



<b>Age of child</b>	<b>Vaccines needed</b>	<b>How and where it is given</b>
At birth	BCG	Right arm
6 weeks	RV (1)	Liquid by mouth
	PCV (1)	Intramuscular Right thigh
	Hexavalent (DTaP-IPV-Hib-HBV) (1)	Intramuscular Left thigh
10 weeks	Hexavalent (DTaP-IPV-Hib-HBV) (2)	Intramuscular Left thigh
14 weeks	Rotavirus (2)	Oral
	PCV (2)	Intramuscular Right thigh
	Hexavalent (DTaP-IPV-Hib-HBV) (3)	Intramuscular Left thigh
6 months	Measles (Not required if giving MMR at 12 months)	Subcutaneous Left thigh
9 months	PCV (3)	Intramuscular Right thigh
12 months	MMR	Subcutaneous Right arm
18 months	Hexavalent (DTaP-IPV-Hib-HBV) (4)	Intramuscular Left arm
6 years	DTaP-IPV	Intramuscular Left arm
12 years	TDaP-IPV	Intramuscular Left arm

#### **Additional Vaccinations**

Girls - 9 years and older	HPV (1) HPV (2)	Intramuscular Non-dominant arm
---------------------------	--------------------	--------------------------------

#### **Abbreviations:**

BCG	Bacilles Calmette Guerin
RV	Rotavirus
DTaP-IPV-Hib-HBV	Diphtheria, Tetanus, Acellular Pertussis, Inactivated Polio Vaccine and Haemophilus Influenzae Type B and Hepatitis B Combined
MMR	Measles, Mumps, Rubella
PCV	Pneumococcal Conjugated Vaccine
TD	Tetanus and reduced strength of Diphtheria Vaccine
DTaP-IPV	Diphtheria, Tetanus, Pertussis, Polio
TDaP-IPV	Tetanus, Diphtheria, Pertussis, Polio
HPV	Human papillomavirus vaccines

